

Long-term multiwavelength studies of high-redshift blazar 0836+710

Akyuz A., Thompson D., Donato D., Perkins J., Fuhrmann L., Angelakis E., Zensus J., Larsson S., Sokolovsky K., Kurtanidze O.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Aims. The observation of γ -ray flares from blazar 0836+710 in 2011, following a period of quiescence, offered an opportunity to study correlated activity at different wavelengths for a high-redshift ($z = 2.218$) active galactic nucleus. **Methods.** Optical and radio monitoring, plus Fermi-LAT γ -ray monitoring provided 2008-2012 coverage, while Swift offered auxiliary optical, ultraviolet, and X-ray information. Other contemporaneous observations were used to construct a broad-band spectral energy distribution. **Results.** There is evidence of correlation but not a measurable lag between the optical and γ -ray flaring emission. In contrast, there is no clear correlation between radio and γ -ray activity, indicating radio emission regions that are unrelated to the parts of the jet that produce the γ rays. The γ -ray energy spectrum is unusual in showing a change of shape from a power law to a curved spectrum when going from the quiescent state to the active state. © ESO, 2013.

<http://dx.doi.org/10.1051/0004-6361/201321721>

Keywords

Galaxies: active, Gamma rays: galaxies, Quasars: individual: 0836+710